

Counterexamples to a conjecture by Gun and co-workers, its correct reformulation and the transcendence of some series

F. M. S. Lima

Institute of Physics, University of Brasilia, P.O. Box 04455, 70919-970, Brasilia, DF, Brazil

Abstract

In a recent work, Gun, Murty and Rath have introduced a ‘theorem’ asserting that $\sum_{n=-\infty}^{\infty} (n + \alpha)^{-k}$ yields a transcendental number for all $\alpha \in \mathbb{Q} \setminus \mathbb{Z}$, k being an integer greater than 1. I show here in this short paper that this conjecture is *false* whenever k is odd and α is a half-integer. I also prove that these are the only counterexamples, which allows for a correct reformulation. The resulting theorem implies the transcendence of both the polygamma function at rational entries and certain zeta series.

Key words: Dirichlet series, Cotangent derivatives, Transcendental numbers, Polygamma function, Zeta series

2000 MSC: 11J81, 11J91, 11M41, 33B15

Email address: `fabio@fis.unb.br` (F. M. S. Lima)

Preprint submitted to : IJMMS

April 8, 2011

1. Introduction

In a recent work on the transcendence of the log-gamma function and some discrete periods, Gun, Murty and Rath (GMR) have presented the following ‘theorem’ on the transcendence of certain Dirichlet series (see Theorem 4.1 in Ref. [1]).

Conjecture 1 (GMR). *Let α be a non-integral rational number and $k > 1$ be a natural number. Then $\sum_{n=-\infty}^{+\infty} \frac{1}{(n+\alpha)^k}$ is a transcendental number.*

In a recent work, I have shown that even the main ‘theorem’ in that work (namely, their Theorem 3.1) has an incorrect proof [2]. So, by suspecting that the above series could converge to an algebraic number for some non-integral rational α , I have considered the possibility of finding a counterexample to their assertion.¹ After some computational tests, I have found a simple counterexample: the series is *null* (hence an algebraic number) for $\alpha = \frac{1}{2}$ and $k = 3$. Of course, this implies that Conjecture 1, above, is *false*.

Here in this work, I show that the original proof by Gun and co-workers is *invalid* when α is a half-integer and k is an odd integer. By repairing their defective proof, I show that these values of α and k compose an (infinite) set containing all possible counterexamples, which allows for a correct reformulation of their assertion. This yields a theorem that determines whether $\sum_{n=-\infty}^{+\infty} 1/(n+\alpha)^k$ is an algebraic or a transcendental number, for all rational α , $\alpha \notin \mathbb{Z}$, and every $k \in \mathbb{Z}$, $k > 1$. My proof is based only upon the periodicity and some basic properties of this series. I also show that

¹The short proof provided in Ref. [1] is, at first sight, quite convincingly.

the theorem has interesting consequences for the transcendence of both the polygamma function at rational entries and certain zeta series.

2. A first simple counterexample

By testing the validity of the GMR conjecture for $k = 3$ and some rational numbers $\alpha \in (0, 1)$, I have found the following counterexample.

Lemma 1 (First counterexample). *The series*

$$\sum_{n=-\infty}^{+\infty} \frac{1}{\left(n + \frac{1}{2}\right)^3} \quad (1)$$

is null, thus an algebraic number.

PROOF. By writing the above series as the sum of two series, one for the non-negative values of n and the other for the negative ones, which is a valid procedure since the series converges absolutely, one has

$$\begin{aligned} \sum_{n=-\infty}^{+\infty} \frac{1}{\left(n + \frac{1}{2}\right)^3} &= \sum_{n=-\infty}^{-1} \frac{1}{\left(n + \frac{1}{2}\right)^3} + \sum_{n=0}^{\infty} \frac{1}{\left(n + \frac{1}{2}\right)^3} \\ &= \sum_{n=-\infty}^{-1} \frac{2^3}{(2n+1)^3} + \sum_{n=0}^{\infty} \frac{2^3}{(2n+1)^3}. \end{aligned}$$

Now, by substituting $n = -m$ in the series for $n < 0$ and $n = j - 1$ in the series for $n \geq 0$, one has

$$\sum_{n=-\infty}^{+\infty} \frac{1}{\left(n + \frac{1}{2}\right)^3} = -8 \sum_{m=1}^{\infty} \frac{1}{(2m-1)^3} + 8 \sum_{j=1}^{\infty} \frac{1}{(2j-1)^3} = 0. \quad (2)$$

□

The existence of a counterexample to the GMR conjecture implies that its original statement (see Theorem 4.1 (2) of Ref. [1]) is *false*. In fact, by scrutinizing the proof furnished there in Ref. [1], I have found some defective points. Their proof reads [1]:

PROOF (GMR *incorrect* PROOF). We know that

$$\sum_{n=-\infty}^{+\infty} \frac{1}{(n+\alpha)^k} = \frac{1}{\alpha} + \frac{(-1)^k}{(k-1)!} D^{k-1}(\pi \cot \pi z) \Big|_{z=\alpha}, \quad (3)$$

where $D := \frac{d}{dz}$. It is a consequence of a result of Okada [3] that $D^{k-1}(\pi \cot \pi z) \Big|_{z=\alpha}$ is non-zero. But then it is π^k times a non-zero linear combination of algebraic numbers of the form $\csc(\pi\alpha)$, $\cot(\pi\alpha)$. Thus we have the result. $\square$

There in the Okada's cited work [3], one finds, in its only theorem, the following (correct) linear independence result.

Lemma 2 (Okada's theorem). *Let k and q be integers with $k > 0$ and $q > 2$. Let T be a set of $\varphi(q)/2$ representatives mod q such that the union $\{T, -T\}$ is a complete set of residues prime to q . Then the real numbers $D^{k-1}(\cot \pi z) \Big|_{z=a/q}$, $a \in T$, are linearly independent over $\mathbb{Q}$.²*

See Ref. [3] for a detailed proof of this lemma based upon the partial fraction decomposition of $D^{k-1}(\pi \cot \pi z)$, valid for all $z \notin \mathbb{Z}$, as well as a theorem by Baker-Birch-Wirsing on cyclotomic polynomials (see his Corollary 1). Note, however, that this lemma says nothing about the cotangent derivatives at $z = a/q$ with $q = 2$. Then, the linear independence over $\mathbb{Q}$ is

²Here, $\varphi(q)$ is the Euler totient function.

not guaranteed if z is a half-integer, which is a source of potential counterexamples to the GMR conjecture. Moreover, the proof of Okada's theorem is based upon the following partial fraction decomposition for his function $F_k(z) = \frac{k}{(-2\pi i)^k} D^{k-1}(\pi \cot \pi z)$, valid for all $z \notin \mathbb{Z}$ (see Eq. (1) of Ref. [3]):

$$-\frac{k!}{(2\pi i)^k} \sum_{n=-\infty}^{+\infty} \frac{1}{(n+z)^k} = \frac{k}{(-2\pi i)^k} D^{k-1}(\pi \cot \pi z) ,$$

which simplifies to

$$\sum_{n=-\infty}^{+\infty} \frac{1}{(n+z)^k} = \frac{(-1)^{k-1}}{(k-1)!} D^{k-1}(\pi \cot \pi z) . \quad (4)$$

This identity, by itself, shows that the expression for $\sum_{n=-\infty}^{+\infty} 1/(n+\alpha)^k$ taken into account by Gun and co-workers in their proof [1], as reproduced in our Eq. (3), is *incorrect*.

3. An infinity of counterexamples

As we are only interested on integer values of k , $k > 1$, let us define the following countable set of real functions:

$$S_k(z) := \sum_{n=-\infty}^{+\infty} \frac{1}{(n+z)^k} , \quad (5)$$

defined for all real $z \notin \mathbb{Z}$.³ From this definition, it is easy to deduce the following mathematical properties.

Lemma 3. *The functions $S_k(z)$, $k \in \mathbb{Z}$, $k > 1$, have the following properties, valid for all z in its domain (i.e., $z \in \mathbb{R} \setminus \mathbb{Z}$):*

³Note that, for all $m \in \mathbb{Z}$, $\lim_{z \rightarrow m} |S_k(z)| = \infty$.

- (i) All functions $S_k(z)$ are periodic, with an unitary period;
- (ii) For even values of k , $S_k(z) > 0$;
- (iii) All functions $S_k(z)$ are differentiable;
- (iv) For odd values of k , $S_k(z)$ are strictly decreasing functions.

PROOF. The property (i) follows from the fact that, $\forall k \in \mathbb{Z}, k > 1$, and for all real $z \notin \mathbb{Z}$, $S_k(z+1) = \sum_{n=-\infty}^{+\infty} 1/[n + (z+1)]^k = \sum_{m=-\infty}^{+\infty} 1/(m+z)^k = S_k(z)$. Note that n has been substituted by $m-1$ in the first series. Property (ii) is an obvious consequence of the fact that, for any positive *even* k , every term $1/(n+z)^k$ of the series that defines $S_k(z)$, see Eq. (5), is positive. Property (iii) follows from the fact that the series for $S_k(z)$ in Eq. (5) is term-by-term differentiable with respect to z , without restrictions for $z \in (0, 1)$, and that differentiation does not affect the convergence for any z in this interval.⁴ This differentiability for $z \in (0, 1)$ can then be extended to all real $z \notin \mathbb{Z}$ — i.e., all points in the domain of $S_k(z)$ — by making use of their periodicity, as established in property (i). Property (iv) follows from a less direct argument. Firstly, from Eq. (4) we deduce that, for any positive integer m ,

$$S_{2m+2}(z) = \frac{-1}{(2m+1)!} D^{2m+1} (\pi \cot \pi z) \quad (6)$$

and

$$\frac{d}{dz} S_{2m+1}(z) = \frac{1}{(2m)!} D^{2m+1} (\pi \cot \pi z). \quad (7)$$

⁴This property also follows from the representation of $S_k(z)$ as a sum/difference of two polygamma functions $\psi_k(z)$ with $z \in (0, 1)$, since each $\psi_k(z)$ is differentiable at all points of this interval. This representation will be considered in the next section.

By isolating the cotangent derivative at the right-hand side of Eq. (7) and then substituting it on Eq. (6), one finds that

$$\frac{d}{dz} S_{2m+1}(z) = -(2m+1) S_{2m+2}(z). \quad (8)$$

By property (ii), $S_{2m+2}(z) > 0$, thus $dS_{2m+1}/dz < 0$ for all real $z \notin \mathbb{Z}$. Then, $S_{2m+1}(z)$ is a strictly decreasing function in all points of its domain. $\square$

An immediate consequence of the periodicity of $S_k(z)$ is the periodic repetition of the null result established for $\alpha = \frac{1}{2}$ in Lemma 1. This leads to an infinite set of counterexamples to the GMR assertion, as establishes the following theorem.

Lemma 4 (More counterexamples for $k = 3$). *For every integer m , the series*

$$\sum_{n=-\infty}^{+\infty} \frac{1}{\left(n + m + \frac{1}{2}\right)^3} \quad (9)$$

is null.

PROOF. From the fact that $S_3\left(\frac{1}{2}\right) = 0$ (see Lemma 1) and from the periodicity of $S_3(z)$, with an unitary period, as established in Lemma 3, one has $S_3\left(m + \frac{1}{2}\right) = 0$.

$\square$

All counterexamples to the GMR original assertion presented hitherto are particular cases of a more general set of counterexamples, as establishes the next lemma.

Lemma 5 (Counterexamples for odd values of k). *For every odd integer k , $k > 1$, and every $m \in \mathbb{Z}$, the series*

$$\sum_{n=-\infty}^{+\infty} \frac{1}{\left(n + m + \frac{1}{2}\right)^k} \quad (10)$$

is null.

PROOF. The proof for $S_k(\frac{1}{2})$ (i.e., for $m = 0$), valid for any odd integer k , $k > 1$, is analogue to that developed in Lemma 1 for $k = 3$. By writing the corresponding series as the sum of two series, one for $n < 0$ and the other for $n \geq 0$, one has

$$\begin{aligned} S_k\left(\frac{1}{2}\right) &= \sum_{n=-\infty}^{+\infty} \frac{1}{\left(n + \frac{1}{2}\right)^k} = \sum_{n=-\infty}^{-1} \frac{1}{\left(n + \frac{1}{2}\right)^k} + \sum_{n=0}^{\infty} \frac{1}{\left(n + \frac{1}{2}\right)^k} \\ &= \sum_{n=-\infty}^{-1} \frac{2^k}{(2n+1)^k} + \sum_{n=0}^{\infty} \frac{2^k}{(2n+1)^k}. \end{aligned}$$

By substituting $n = -m$ in the series for $n < 0$ and $n = j - 1$ in the series for $n \geq 0$, one finds the following null result:

$$S_k\left(\frac{1}{2}\right) = -2^k \sum_{m=1}^{\infty} \frac{1}{(2m-1)^k} + 2^k \sum_{j=1}^{\infty} \frac{1}{(2j-1)^k} = 0. \quad (11)$$

The extension of this null result to all other half-integer values of z follows from the fact that $S_k(z)$ is periodic, with an unitary period (see property (i) of Lemma 3). Then, $S_k\left(m + \frac{1}{2}\right) = 0$.

□

With these properties and counterexamples in hands, we can now reformulate the GMR conjecture.

4. Reformulating the GMR conjecture

Let us present and prove a theorem which determines whether $S_k(\alpha)$ is an algebraic number or not for non-integral rational values of α .

Theorem 1 (Main result). *For any integer k , $k > 1$, and every $\alpha \in \mathbb{Q} \setminus \mathbb{Z}$, the series*

$$\sum_{n=-\infty}^{+\infty} \frac{1}{(n + \alpha)^k} \quad (12)$$

*is either an algebraic multiple of π^k or null. It is null **if and only if** k is an odd integer and α is a half-integer.*

PROOF. From Eqs. (4) and (5), we know that, for any integer k , $k > 1$,

$$S_k(\alpha) = \sum_{n=-\infty}^{+\infty} \frac{1}{(n + \alpha)^k} = \frac{(-1)^{k-1}}{(k-1)!} D^{k-1} (\pi \cot \pi z) \Big|_{z=\alpha}. \quad (13)$$

From Okada's theorem, as reproduced in Lemma 2, one knows that, for all rational $\alpha \notin \mathbb{Z}$ and any integer k , $k > 1$, $D^{k-1} (\pi \cot \pi z) \Big|_{z=\alpha}$ is *non-zero*, the only possible exceptions being the half-integer values of α (i.e., non-integer fractions with a denominator equal to 2). From the usual rules for derivatives, it is easy to note that when the above cotangent derivative is non-zero, it is π^k times a non-zero linear combination (with integer coefficients) of algebraic numbers of the form $\csc(\pi \alpha)$ and $\cot(\pi \alpha)$. When that derivative is null, the resulting equation $S_k(\alpha) = 0$ has no real roots if k is even, according to property (ii) of Lemma 3. For odd values of k , on the other hand, all half-integer values of α are roots of $S_k(\alpha) = 0$, as establishes Lemma 5.

All that rests is to prove that the half-integers are the *only* roots of $S_{2\ell+1}(\alpha) = 0$, ℓ being any positive integer. For this, let us restrict our

attention to the open interval $(0, 1)$. Note that $\alpha = \frac{1}{2}$ belongs to the interval $(0, 1)$ and is a root of $S_{2\ell+1}(\alpha) = 0$ for all positive integer ℓ , as guaranteed by Lemma 5. From properties (iii) and (iv) of Lemma 3, we know that $S_{2\ell+1}(\alpha)$ is a *strictly decreasing* differentiable function, for all real $\alpha \in (0, 1)$. Then, $S_{2\ell+1}(\alpha_1) \neq S_{2\ell+1}(\alpha_2)$ for all distinct $\alpha_1, \alpha_2 \in (0, 1)$. In particular, $S_{2\ell+1}(\alpha)$ cannot be null for two distinct values of its argument, both belonging to $(0, 1)$. It follows that there is at most one real root in $(0, 1)$. Therefore, $\alpha = \frac{1}{2}$ is the *only* root in the interval $(0, 1)$. Finally, the periodicity of $S_{2\ell+1}(\alpha)$ guarantees that $\alpha = \frac{1}{2} + m$ (i.e., the half-integers) are the only real solutions for $S_{2\ell+1}(\alpha) = 0$, $\alpha \notin \mathbb{Z}$.

□

From the fact that π is a transcendental number, as first proved by Lindemann (1882), it follows that

Corollary 1 (Reformulation of the GMR conjecture). *For any integer k , $k > 1$, and every $\alpha \in \mathbb{Q} \setminus \mathbb{Z}$, the series*

$$\sum_{n=-\infty}^{+\infty} \frac{1}{(n + \alpha)^k} \tag{14}$$

*is either null or a transcendental number. It is null **if and only if** k is an odd integer and α is a half-integer.*

Another interesting consequence of Theorem 1 comes from the fact that we can easily write the cotangent derivatives in Eq. (4) in terms of the polygamma function $\psi_k(z) := \psi^{(k)}(z) = d^k \psi(z)/dz^k$, where $\psi(z) := \frac{d}{dz} \ln \Gamma(z)$

is the so-called digamma function. From the reflection formula for $\psi_k(z)$, namely [4, 5]

$$\psi_k(1-z) - (-1)^k \psi_k(z) = (-1)^k D^k(\pi \cot \pi z), \quad (15)$$

valid for all non-negative integers k , with $\psi_0(z) := \psi(z)$, and $z \notin \mathbb{Z}$, and taking into account Eqs. (4) and (5), one finds

$$S_k(\alpha) = \frac{\psi_{k-1}(1-\alpha) + (-1)^k \psi_{k-1}(\alpha)}{(k-1)!}. \quad (16)$$

From Theorem 1, we know that, for any integer $k > 1$ and every $\alpha \in \mathbb{Q} \setminus \mathbb{Z}$, $S_k(\alpha)$ is either null or an algebraic multiple of π^k . Of course, the same conclusion is valid for $(k-1)! S_k(\alpha)$. By taking Eq. (16) into account, we have the following result.

Corollary 2 (Transcendence of the polygamma function). *For any positive integer k and every $\alpha \in \mathbb{Q} \setminus \mathbb{Z}$,*

$$\psi_k(1-\alpha) - (-1)^k \psi_k(\alpha) \quad (17)$$

*is either null or an algebraic multiple of π^{k+1} . It is null **if and only if** α is a half-integer and k is even.*

Finally, let us make use of the above corollaries for establishing a result on the algebraic nature of certain zeta series. In Eq. (7) of Sec. 1.41 of Ref. [6], one finds the Taylor series expansion

$$z \cot z = \sum_{n=0}^{\infty} (-1)^n \frac{2^{2n} B_{2n}}{(2n)!} z^{2n}, \quad (18)$$

which converges for all real z with $|z| < \pi$. By exchanging z by πz and making use of the Euler's formula for even zeta values, namely

$$\zeta(2n) = (-1)^{n-1} \frac{2^{2n-1} B_{2n} \pi^{2n}}{(2n)!}, \quad (19)$$

where $\zeta(m) := \sum_{n=1}^{\infty} 1/n^m$ is the Riemann zeta function, it is easy to deduce that

$$\pi \cot(\pi z) = 1/z - 2 \sum_{n=1}^{\infty} \zeta(2n) z^{2n-1}, \quad (20)$$

with the series at the right-hand being convergent for all real z with $|z| < 1$.⁵ From this equation, by calculating the successive derivatives on both sides, it is easy to derive the following general formulae for the derivatives of order m , m being any non-negative integer:

$$D^{2m}(\pi \cot \pi z) = \frac{(2m)!}{z^{2m+1}} - 2 \sum_{n=m+1}^{\infty} \zeta(2n) \cdot (2n-1) \cdot \dots \cdot (2n-2m) z^{2n-2m-1} \quad (21)$$

and

$$D^{2m+1}(\pi \cot \pi z) = -\frac{(2m+1)!}{z^{2m+2}} - 2 \sum_{n=m+1}^{\infty} \zeta(2n) \cdot (2n-1) \cdot \dots \cdot (2n-2m-1) z^{2n-2m-2} \quad (22)$$

From Eq. (4) and Corollary 1, one readily deduces that

Corollary 3 (Transcendence of certain zeta series). *For any positive integer m and every rational $z \in (-1, 1) \setminus \{0\}$, both the zeta series*

$$\sum_{n=m+1}^{\infty} \zeta(2n) \cdot (2n-1) \cdot \dots \cdot (2n-2m) z^{2n-2m-1} \quad (23)$$

⁵For $z = 0$, the series is null, but Eq. (20) is not valid due to an obvious division by zero.

and

$$\sum_{n=m+1}^{\infty} \zeta(2n) \cdot (2n-1) \cdot \dots \cdot (2n-2m-1) z^{2n-2m-2} \quad (24)$$

converge to some algebraic multiple of π^{2m+1} and π^{2m+2} , respectively, the only exceptions being for the former series with $z = \pm \frac{1}{2}$, for which it is null.

Acknowledgments

The author acknowledges a postdoctoral fellowship from CNPq (Brazilian agency) during the course of this work.

References

- [1] S. Gun, M. R. Murty, P. Rath, *Transcendence of the log gamma function and some discrete periods*, J. Number Theory **129** (2009) 2154–2165.
- [2] F. M. S. Lima, *On the possible exceptions for the transcendence of the log-gamma function at rational entries*. ArXiv: 0908.3253v2 (2009).
- [3] T. Okada, *On an extension of a theorem of S. Chowla*, Acta Arith. **38** (1980/1981) 341–345.
- [4] K. S. Kölbig, *The polygamma function and the derivatives of the cotangent function for rational arguments*, CERN Report CN/96/5 (1996).
- [5] E. W. Weisstein, *Polygamma Function*. From *MathWorld*. Available at:
 $\backslash\protect\vrule\width0pt\protect\href{\texttt{http://mathworld.wolfram.com/PolygammaFunction.html}}$
- [6] I. S. Gradshteyn and I. M. Ryzik, *Table of Integrals, Series, and Products*, 7th ed., Academic Press, New York, 2007.